

An Observational Study of Negative and Low Pressure Hydrocephalus: The Diagnosis and Treatment Jian Yu MD

Department of neurosurgery, Huashan hospital Fudan University

### Introduction

Negative or low pressure hydrocephalus, a very rare condition of unknown etiology and optimal treatment, usually present postneurosurgery with clinical and imaging features of hydrocephalus, but with negative or low cerebrospinal fluid pressure. It's important to understand the characteristics of low or negative hydrocephalus and correct management strategy, which are the keys to well results for these patients.

# Methods

11 patients of negative and low pressure hydrocephalus were prospectively collected in this study. The initial intracranial pressure of all patients included in this study was lower than 6 cmH2O. A standard protocol for the treatment of negative and low pressure hydrocephalus is used for the treatment of these

patients, including continuous intracranial pressure monitoring with horizontal positioning, head down and legs elevated to 10-15 degrees, neck wrapping, chest and abdomen bandages, infusion of 5% dextrose fluid, daily cerebrospinal fluid drainage and so on. Daily intracranial pressure, CSF drainage volumes and simultaneous CT images variation are collected. The CSF

collected from the drainage tube was examined daily to rule out infection secondary to prolonged EVD. GOS-E scores were followed up three months later, and management strategies were discussed in detail

## Results

The median of initial intracranial pressure was 2 (interquartile range, -3 to 6) cm H2O. The intracranial pressure was successfully treated to achieve 8 cm H2O (interquartile range, 6 to 14) under continuous intracranial pressure monitoring with horizontal

positioning, head down and legs elevated to 10-15 degrees, neck wrapping for controlled venous drainage, chest and abdomen bandages, infusion of 5% dextrose fluid to lower plasma osmolarity (Na+, 130-135 mmol/L), daily cerebrospinal fluid drainage >200 mL, and arterial blood gas partial pressure of carbon dioxide >40 mm Hg.

# References

1. Cohen AR. Editorial: Low-pressure hydrocephalus. Journal of Neurosurgery. 2011;115(5):1029-1030

2. Pang D, Altschuler E. Low-Pressure Hydrocephalic State and Viscoelastic Alterations in the Brain. Neurosurgery. 1994;35(4):643-656.

3. Smalley ZS, Venable GT, Einhaus S, Klimo Jr P. Low-pressure Hydrocephalus in Children: A Case Series and Review of the Literature. Neurosurgery. 2016;80(3):439-447.

4. Lin B-J, Ma H-I. A rare case of low-pressure hydrocephalus after skull base surgery. Neurosurgery Quarterly. 2016;26(3):256-258.

5. Hunn BH, Mujic A, Sher I, Dubey AK, Peters-Willke J, Hunn AW. Successful treatment of negative pressure hydrocephalus using timely titrated external ventricular drainage: a case series. Clin Neurol Neurosurg. Jan 2014;116:67-71.

6. Dandy WE. Intracranial tumors and abscesses causing communicating hydrocephalus. Annals of surgery. 1925;82(2):199.

7. Akins PT, Guppy KH, Axelrod YV, Chakrabarti I, Silverthorn J, Williams AR. The genesis of low pressure hydrocephalus. Neurocrit Care. Dec 2011;15(3):461-468.

### **Learning Objectives**

Better diagnosis and treatment of negative and low pressure hydrocephalus.

#### Conclusions

Negative and low pressure hydrocephalus are a complex condition with an unknown etiology and equally challenging treatment strategy. It illustrates that titration EVD should be combined with other therapies to restore positive pressure under continuous ICP

monitoring. Furthermore, mild hyponatremia (130 mmol/L) and supine position should be continued even after titration EVD is suspended.